

A SOLUTION TO THE L SPACE PROBLEM AND RELATED ZFC CONSTRUCTIONS

JUSTIN TATCH MOORE

ABSTRACT. In this paper I will construct a non-separable hereditarily Lindelöf space (L space) without any additional axiomatic assumptions. I will also show that there is a function $f : [\omega_1]^2 \rightarrow \omega_1$ such that if $A, B \subseteq \omega_1$ are uncountable and $\xi < \omega_1$, then there are $\alpha < \beta$ in A and B respectively with $f(\alpha, \beta) = \xi$. Previously it was unknown whether such a function existed even if ω_1 was replaced by 2. Finally, I will prove that there is no basis for the uncountable regular Hausdorff spaces of cardinality $\aleph_1$. Each of these results gives a strong refutation of a well known and longstanding conjecture. The results all stem from the analysis of oscillations of coherent sequences $\langle e_\alpha : \alpha < \omega_1 \rangle$ of finite-to-one functions. I expect that the methods presented will have other applications as well.

1. INTRODUCTIONS

One goal in mathematics is to classify structures, particularly those which lie at the core of mathematics — graphs, groups, orders, relations, solids, surfaces, vector spaces, and topologies.¹ Classifications can come in many forms. The strongest and most infrequent is a complete and informative listing — up to presentation — of the structures of a given class of objects. The following are examples of such a classification.

2000 *Mathematics Subject Classification*. Primary: 54D20, 54D65, 03E02, 03E75.

Key words and phrases. L space, negative partition relation, Tukey order, hereditarily Lindelöf, hereditarily separable, basis.

This paper is dedicated to Stevo Todorčević for teaching me how to traverse ω_1 and for his inspirational [28]. I would like to thank Boban Veličković for his careful reading of early drafts of this paper. The clarity of the present arguments owe much to Todorčević's and Veličković's suggestions. A number of other people — too many to name — have also made the generous contribution of reading the paper and offering suggestions. The research presented in this paper was funded by NSF grant DMS-0401893.

¹All topological spaces in this paper are regular and Hausdorff.

Theorem 1.1. *There are five regular polyhedra: the tetrahedron, the cube, the octahedron, the dodecahedron, and the icosahedron.*

Theorem 1.2. *Any two closed orientable 2-manifolds with the same genus are homeomorphic.*

Theorem 1.3. [3] *Any countable transitive relation is Tukey equivalent to one of the following forms for some non-negative integers n_i ($i < 3$):*

$$n_0 \cdot 1 \oplus n_1 \cdot \omega \oplus n_2 \cdot \omega^{<\omega}.$$

Frequently it is not possible to have a reasonable complete classification of this sort. If this happens, it is still possible to ask whether the objects can be classified from “above” or from “below.” In each case, there is a need for a notion of reduction $\leq$ (e.g. being a minor, a normal subgroup, subgraph, a subspace, etc.). A family $\mathcal{B}$ is a *basis* for a class $\mathcal{C}$ if for any member C of $\mathcal{C}$, there is a B in $\mathcal{B}$ such that $B \leq C$. A family $\mathcal{U}$ is a *universal* for $\mathcal{C}$ if any member C of $\mathcal{C}$, there is a U in $\mathcal{U}$ such that $C \leq U$. An example of a basis theorem is the following result of Kuratowski.

Theorem 1.4. [15] *If G is a non-planar graph, then G contains either $K_{3,3}$ or K_5 as a minor.*

The classification of finite simple groups can also be considered a basis theorem for the class of finite groups with reduction being “is contained as a normal subgroup.” The following are two examples of universality results.

Theorem 1.5. $\mathbb{R}$ *is a universal separable linear order.*

Theorem 1.6. *The irrationals are surjectively universal for the separable completely metrizable spaces.*²

In this paper we will focus on basis problems. Since the whole class $\mathcal{C}$ is trivially a basis, the goal is to find a basis which is informative. This is often quantified in terms of restrictions on its cardinality. In cases where the basis must be infinite, it is also of interest to know whether it can contain only minimal elements, hence avoiding any redundancies.

In some cases it is possible to show that there is no reasonable basis — that any such basis is either large in terms of cardinality or must contain redundancies. While such a result is not exactly in line with the goal of classification, it is still informative as to the limits of our ability to understand the class.

²Unfortunately it is not even well known in many mathematical circles that the irrationals are themselves completely metrizable.

A phenomenon of this sort has been repeatedly observed for objects at the level of the continuum. The following classical result of Sierpiński is a typical example.

Theorem 1.7. [24] *There is a subset $X \subseteq \mathbb{R}$ of cardinality continuum such that if $f \subseteq X^2$ is a continuous injection, then f differs from the identity map on a set of size less than continuum. In particular, the suborders of X witness that any basis for the separable linear orders of cardinality continuum must be uncountable and contain redundancies.*

In this paper we will focus on basis problems for uncountable objects. For finite and countable objects, classification theorems are typically decidable by the accepted axioms of mathematics (ZFC). This phenomenon is largely explained by Shoenfield’s absoluteness theorem [23]. When dealing with uncountable objects, however, it is often the case that additional axioms are necessary if any classification is consistent with ZFC. This is because the non-existence of reductions between fixed mathematical objects can be fail to be absolute. Axioms such as Jensen’s $\diamond$ and the Continuum Hypothesis allow one to inductively define examples while diagonalizing against a list of all possible reductions. If $2^{\aleph_0} = \aleph_1$, then the pathologies mentioned above occur already at the first uncountable cardinal. For instance, Sierpiński’s theorem easily yields the following fact.

Theorem 1.8. *If the Continuum Hypothesis is true, then any basis for the uncountable linear or the uncountable topological spaces must have cardinality at least $2^{\aleph_1}$ and contain redundancies.*

On the other hand, the lack of embeddings between two uncountable sets of reals is not absolute as the following result of Baumgartner’s demonstrates.

Theorem 1.9. [2] *If X and Y are two uncountable sets of reals, then there is a forcing which preserves uncountability and which adds an order preserving map from X into Y . Moreover, the Proper Forcing Axiom implies that such an embedding already exists provided X has cardinality $\aleph_1$.*

The Proper Forcing Axiom (PFA) is a statement which allows one to find embeddings *generically* between two objects of cardinality $\aleph_1$. The use of this axiom can be likened to that of Erdős’s probabilistic method in finite combinatorics (see [1]). The main difference is that the notion of a probability space is abstracted and an axiom is required for the deduction “if an object can be selected with positive probability, then it exists.” This axiom is natural to assume if one wishes to classify

objects at the level of $\aleph_1$, the smallest uncountable cardinal. In fact, there is the following empirical fact:³ For a given class $\mathcal{C}$ consisting of structures of size $\aleph_1$ the existence of a classification or a basis is either a consequence of PFA or else is provably false without additional axiomatic assumptions. This has been partially explained by the work of Woodin [36].

In his 1998 address to the International Congress of Mathematicians, Todorćević outlined a number of basis conjectures⁴ — in the presence of PFA — which had been under consideration since the 1960's and 70's.

Conjecture 1. (*Todorćević; [28]*) *If R is a binary relation, then either $R \leq \aleph_0 \cdot \omega_1$ or $[\omega_1]^{<\aleph_0} \leq R$.*

Conjecture 2. (*Hajnal, Juhasz; [9]*) *If X is a regular Hausdorff space, then the following are equivalent:*

- (1) *X is hereditarily separable.*
- (2) *X is hereditarily Lindelöf.*
- (3) *X does not contain an uncountable discrete subspace.*

Conjecture 3. (*Galvin⁵*) *The partition relation $\omega_1 \rightarrow (\omega_1; \omega_1)_2^2$ holds.*

Conjecture 4. (*Shelah; [22]*) *The uncountable linear orders have a five element basis consisting of X , ω_1 , ω_1^* , C , and C^* where X is a set of reals of cardinality $\aleph_1$ and C is a Countryman type.⁶*

Conjecture 5. (*Gruenhage; [8]*) *The uncountable regular Hausdorff spaces have a three element basis consisting of a set of reals of cardinality $\aleph_1$ with the metric, the Sorgenfrey, and the discrete topology.*

Conjecture 6. (*Fremlin; DN of [5]*) *If X is a compact space in which closed sets are G_δ , then there is an at most two-to-one map from X into a compact metric space.*

Todorćević has, assuming PFA, proved that Conjecture 1 for the transitive relations [32], and confirmed the equivalence of (1) and (3) in Hajnal and Juhasz's problem [27]. In [18], I confirmed that Conjecture 4 follows from PFA. Gruenhage has proved that Conjecture 5 holds

³As with any claim of this sort, there are counterexamples. A notable example is Katětov's problem — see [17].

⁴Todorćević lists these as conjectures in [28]. This is not to say that the author of the statement necessarily speculated a certain resolution.

⁵This attribution is a conjecture of Hajnal. The first time this appeared in print seems to be [4].

⁶A linear order C is Countryman if it is uncountable and yet C^2 is the union of countably many chains in the coordinate-wise order.

for the class of cometrizable spaces assuming PFA [7]. Todorćević has verified that Conjecture 6 holds for the Rosenthal compacta in ZFC [29].

In this paper I will prove results in the other direction, refuting all of the conjectures except Conjectures 4 and 6.

Theorem 1.10. *There is a function $f : [\omega_1]^2 \rightarrow \omega_1$ such that whenever $A, B \subseteq \omega_1$ are uncountable and $\xi < \omega_1$ there are α in A and β in B such that $\alpha < \beta$ and $f(\alpha, \beta) = \xi$. In particular $\omega_1 \not\rightarrow (\omega_1; \omega_1)_2^2$.*

Theorem 1.11. *There is a family $\mathcal{F}$ of binary relations which contains an antichain of cardinality $2^{\aleph_1}$, is downwards directed and such that for all R in $\mathcal{F}$, $R \not\leq \aleph_0 \cdot \omega_1$ and $[\omega_1]^{<\aleph_0} \not\leq R$.*

Theorem 1.12. *There is a non-separable, hereditarily Lindelöf topological space.*

Theorem 1.13. *Any basis for the uncountable topological spaces has cardinality strictly greater than $\aleph_1$.*

It is worth noting that the existence of an L space alone gives a refutation of these conjectures. The above results, however, take the failures of these conjectures a step further and require slightly different combinatorial refinements.

The theorems in this paper are a consequences of the following result and its technical strengthenings. The reader is referred to Section 2 for the undefined notions.

Theorem 1.14. *Suppose that $\langle e_\beta : \beta < \omega_1 \rangle$ is a coherent sequence of finite-to-one functions with $e_\beta : \beta \rightarrow \omega$ for each $\beta < \omega_1$ and suppose that $L : [\omega_1]^2 \rightarrow [\omega_1]^{<\aleph_0}$ satisfies the properties of a lower trace function. If $A, B \subseteq \omega_1$ are uncountable, then the set of integers of the form*

$$\text{osc}(e_\alpha \upharpoonright L(\alpha, \beta), e_\beta \upharpoonright L(\alpha, \beta))$$

for $\alpha < \beta$ with α in A , β in B contains arbitrarily long intervals.

This can be likened to the following result of Todorćević, which he used to draw a number of conclusions about Conjecture 2.

Theorem 1.15. [31] *If X and Y are unbounded and countably directed in $(\omega^\omega, <^*)$ and consist of monotonic functions, then there is a natural number l such that for all n there are x and y in X and Y respectively such that*

$$\text{osc}(x, y) = n + l.$$

The paper is organized as follows. Section 2 provides some background on the method of minimal walks and introduces the lower trace function which is used in the statement of the main technical theorem on oscillations. Section 3 provides a review of elementary submodels. The main technical results of the paper are proved in Section 4. Theorem 1.10 is then deduced in Section 5 and a coloring $o : [\omega_1]^2 \rightarrow \omega$ is introduced which harnesses most of the strength of the more technical main theorem. Theorem 1.11 is then deduced in Section 6 after some motivation is provided. The paper closes with Section 7 where Theorems 1.12 and 1.13 are proved. This final section also contains a basic analysis of the L space which is constructed.

This paper is intended to be accessible to any interested reader who is fluent in set theory — basic background can be found in, e.g., [11], [14]. Elementary submodels will be employed in a number of points in the argument. Ironically this represents the only non-elementary technique used in the proof. The essentials are reviewed in Section 3. The reader is referred to III.1 of [10] for more information on elementary submodels and stationary sets. The proofs will also employ the method of minimal walks introduced in [30]. The necessary background is presented in Section 2. The reader is referred to [33] for further reading on minimal walks.

The notation is fairly standard. All ordinals are von Neumann ordinals — they are the set consisting of their predecessors. In particular, $n = \{0, 1, \dots, n-1\}$ and the first infinite ordinal ω is the set of all finite ordinals $\{0, 1, 2, \dots\}$ and is identified with the natural numbers. All counting starts at 0. If k is a natural number and X is a set, then $[X]^k$ is the set of all k -element subsets of X . If X has a canonical linear ordering associated with it and a is in $[X]^k$, then a will be identified with the increasing sequence which enumerates it. If a and b are finite subsets of ω_1 , then $a < b$ will be used to abbreviate “ $\alpha < \beta$ whenever α is in a and β is in b .” Similarly one defines statements such as $\alpha < b$ and $a < \beta$ if α and β are ordinals.

2. THE TRACE FUNCTIONS

In this section I will provide the necessary background on minimal walks. Minimal walks are facilitated by a *C-sequence* or ladder system which one uses to “walk” from an ordinal β down to a smaller ordinal α . Here a *C-sequence* is a sequence $\langle C_\alpha : \alpha < \omega_1 \rangle$ such that C_α is a cofinal subset of α and if $\gamma < \alpha$ then $C_\alpha \cap \gamma$ is finite. It will be useful at certain points to assume that 0 is an element of every C_α .

The following two functions will be of interest to us. The upper trace will not be necessary, but is useful in making the other definitions more transparent.

Definition. [33] (*upper trace*) If $\alpha < \beta$, then $\text{Tr}(\alpha, \beta)$ is defined recursively by

$$\begin{aligned}\text{Tr}(\alpha, \alpha) &= \emptyset, \\ \text{Tr}(\alpha, \beta) &= \text{Tr}(\alpha, \min(C_\beta \setminus \alpha)) \cup \{\beta\}.\end{aligned}$$

Definition. (*lower trace*) If $\alpha < \beta$, then $L(\alpha, \beta)$ is defined recursively by

$$\begin{aligned}L(\alpha, \alpha) &= \emptyset, \\ L(\alpha, \beta) &= L(\alpha, \min(C_\beta \setminus \alpha)) \cup \{\max(C_\beta \cap \alpha)\} \setminus \max(C_\beta \cap \alpha).\end{aligned}$$

Hence the upper trace $\text{Tr}(\alpha, \beta)$ is enumerated by the sequence $\beta = \beta_0 > \beta_1 > \dots > \beta_{l-1}$ where $\beta_{i+1} = \min(C_{\beta_i} \setminus \alpha)$ and $\alpha = \beta_l$ is in $C_{\beta_{l-1}}$. It is easily verified that the lower trace $L(\alpha, \beta)$ is listed as $\xi_0 \leq \xi_1 \leq \dots \leq \xi_{l-1}$ where ξ_i is the maximum of

$$\bigcup_{j=0}^i C_{\beta_j} \cap \alpha.$$

While the upper trace is well studied, this is, to my knowledge, the first analysis of the lower trace (the *full lower trace* $F(\alpha, \beta)$ is a different matter — see [33]).

The lower trace can be axiomatized by the following facts.

Fact 1. If $\alpha \leq \beta \leq \gamma$ and $L(\beta, \gamma) < L(\alpha, \beta)$, then

$$L(\alpha, \gamma) = L(\alpha, \beta) \cup L(\beta, \gamma).$$

Proof. Let α , β , and γ satisfy the hypothesis of the fact. Observe that $L(\beta, \gamma) < \alpha$ and hence $C_\zeta \cap \alpha = C_\zeta \cap \beta$ whenever ζ is in $\text{Tr}(\beta, \gamma)$. Hence β is in $\text{Tr}(\alpha, \gamma)$ and $\text{Tr}(\alpha, \gamma) = \text{Tr}(\alpha, \beta) \cup \text{Tr}(\beta, \gamma)$. Let

$$\gamma = \gamma_0 > \gamma_1 > \dots > \gamma_{l-1}$$

enumerate $\text{Tr}(\alpha, \gamma)$ and let l_0 be such that $\beta = \gamma_{l_0}$. Then $L(\alpha, \gamma)$ is listed as

$$\xi_0 \leq \xi_1 \leq \dots \leq \xi_{l-1}$$

where

$$\xi_i = \max \bigcup_{j=0}^i C_{\gamma_j} \cap \alpha.$$

If $j < l_0$, then $C_{\gamma_j} \cap \alpha = C_{\gamma_j} \cap \beta$ and so $L(\beta, \gamma) = \{\xi_j\}_{j=0}^{l-1}$. On the other hand,

$$\max(C_{\gamma_{l_0}} \cap \alpha) > \xi_{l_0-1}$$

and so

$$\max \bigcup_{j=0}^i (C_{\gamma_j} \cap \alpha) = \max \bigcup_{j=l_0}^i (C_{\gamma_j} \cap \alpha)$$

and hence $L(\alpha, \beta) = \{\xi_j\}_{j=l_0}^{l-1}$. $\square$

Fact 2. *If δ is a limit ordinal, then*

$$\lim_{\xi \rightarrow \delta} \min L(\xi, \delta) = \delta.$$

Proof. This follows immediately from the observation that

$$\min L(\xi, \delta) = \max(C_\delta \cap \xi).$$

$\square$

For the stronger coloring o used later in the paper, we will also need a fixed enumeration $\langle w_\xi : \xi < \omega_1 \rangle$ of $C(2^\omega, \omega)$ which lists all elements stationarily often. Here $C(2^\omega, \omega)$ is the set of all continuous functions from 2^ω into ω . Since 2^ω is compact and ω is discrete, this collection is countable and its elements can be thought of as weighted clopen partitions of 2^ω . We will also need a fixed sequence z_α ($\alpha < \omega_1$) of distinct elements of 2^ω .

This allows for the following definition.

Definition. *If $\alpha < \beta$, then $\mu(\alpha, \beta)$ is the function defined on $L(\alpha, \beta)$ recursively by*

$$\begin{aligned} \mu(\alpha, \beta)(\max(C_\beta \cap \alpha)) &= w_\beta, \\ \mu(\alpha, \beta)(\gamma) &= \mu(\alpha, \min(C_\beta \setminus \alpha))(\gamma) \end{aligned}$$

for all $\gamma > \max(C_\beta \cap \alpha)$ in $L(\alpha, \beta)$. Let $\mu(\alpha, \beta; \xi)$ be the pointwise evaluation of $\mu(\alpha, \beta)$ at z_ξ .

It can be checked that $\mu(\alpha, \beta)$ is the mapping defined by $\xi_i \mapsto w_{\beta_i}$ if $\xi_{i-1} < \xi_i$ or $i = 0$ where ξ_i and β_i are as above. Like L , the function μ can be axiomatized by the following properties.

Fact 3. *If $\alpha < \beta < \gamma$ and $L(\beta, \gamma) < L(\alpha, \beta)$, then*

$$\mu(\alpha, \gamma) = \mu(\alpha, \beta) \cup \mu(\beta, \gamma)$$

(i.e. $\mu(\alpha, \beta)$ and $\mu(\beta, \gamma)$ are both restriction of $\mu(\alpha, \gamma)$).

Fact 4. *If $\xi < \delta$, then $\mu(\xi, \delta)(\min L(\xi, \delta)) = w_\delta$.*

The main technical result of the paper will be concerned with counting oscillations between elements of a coherent⁷ sequence $\langle e_\beta : \beta < \omega_1 \rangle$

⁷A family of functions is *coherent* in this context if any two differ on a finite set on the intersection of their domains.

where e_β is a finite-to-one function from β into ω . The following construction gives a standard example of a coherent family of finite-to-one functions.

Definition. [30] (*maximal weight*) If $\alpha \leq \beta$, then $\varrho_1(\alpha, \beta)$ is defined recursively by

$$\begin{aligned}\varrho_1(\alpha, \alpha) &= 0 \\ \varrho_1(\alpha, \beta) &= \max \left(|C_\beta \cap \alpha|, \varrho_1(\alpha, \min(C_\beta \setminus \alpha)) \right).\end{aligned}$$

Alternately, $\varrho_1(\alpha, \beta)$ is the maximum value of the form $|C_\xi \cap \alpha|$ where ξ ranges over $\text{Tr}(\alpha, \beta)$. Put $e_\beta(\alpha) = \varrho_1(\alpha, \beta)$.

Fact 5. [30] $\langle e_\beta : \beta < \omega_1 \rangle$ is a coherent sequence of finite-to-one functions.

Proof. Let $\beta \leq \beta' < \omega_1$ and $n < \omega$ be given and set D equal to the set of all $\alpha < \beta$ such that either $e_\beta(\alpha) \leq n$ or $e_\beta(\alpha) \neq e_{\beta'}(\alpha)$. It suffices to show that D has no limit points.

To this end, suppose $\delta \leq \beta$. It is easy to check that there is a $\delta_0 < \delta$ such that

$$\begin{aligned}\text{Tr}(\alpha, \beta) &= \text{Tr}(\alpha, \delta) \cup \text{Tr}(\delta, \beta), \\ \text{Tr}(\alpha, \beta') &= \text{Tr}(\alpha, \delta) \cup \text{Tr}(\delta, \beta'), \\ |C_\delta \cap \alpha| &> n\end{aligned}$$

whenever $\delta_0 < \alpha < \delta$. If $\delta_0 < \alpha < \delta$, then

$$\begin{aligned}e_\beta(\alpha) &\geq |C_\delta \cap \alpha| > n \\ e_\beta(\alpha) &= e_\delta(\alpha) = e_{\beta'}(\delta)\end{aligned}$$

and hence α is not in D . Consequently δ is not a limit point of D . $\square$

Remark. It is interesting to note that while coherence and the finite-to-one property are at tension with each other, the verifications of these properties in the previous fact are virtually identical.

Throughout this paper we will now assume that L , μ , $\langle e_\beta : \beta < \omega_1 \rangle$, $\langle w_\alpha : \alpha < \omega_1 \rangle$, and $\langle z_\alpha : \alpha < \omega_1 \rangle$ are fixed objects defined as above.

3. BASIC FACTS ABOUT ELEMENTARY SUBMODELS

In the proof of the main theorem, we will need the following facts about elementary submodels of $H(\aleph_2)$. Recall that $H(\aleph_2)$ is the collection of all sets of hereditary cardinality less than $\aleph_2$. This structure is of interest to us since it contains all of the objects relevant to us and satisfies all of the axioms of ZFC except the power set axiom. An

elementary submodel M of $H(\aleph_2)$ is a subset of $H(\aleph_2)$ such that whenever ϕ is a logical formula with parameters in M , M satisfies ϕ if and only if $H(\aleph_2)$ satisfies ϕ . The following fact is well known.

Fact 6. *If M is a countable elementary submodel of $H(\aleph_2)$, then $M \cap \omega_1$ is an ordinal. Moreover, if F is a finite subset of $H(\aleph_2)$, then the set of all ordinals of the form $M \cap \omega_1$ such that M is a countable elementary submodel of $H(\aleph_2)$ contains a club.⁸*

The following standard facts are very useful when working with countable elementary submodels.

Fact 7. *If M is a countable elementary submodel of $H(\aleph_2)$ which contains some element X , then X is countable iff $X \subseteq M$.*

Fact 8. *If M is an elementary submodel of $H(\aleph_2)$ which contains as an element some subset A of ω_1 , then A is uncountable iff $A \cap M \cap \omega_1$ is unbounded in $M \cap \omega_1$.*

Fact 9. *If M is an elementary submodel of $H(\aleph_2)$, X is in $H(\aleph_2)$ and X is definable from a formula with parameters in M , then X is in M .*

4. OSCILLATIONS ON THE LOWER TRACE

In this section I will prove main technical lemmas of the paper. We will take the following as our definition of the oscillation map.

Definition. *Suppose that s and t are two functions defined on a common finite set of ordinals F . Let $\text{Osc}(s, t; F)$ be the set of all ξ in F such that $s(\xi^-) \leq t(\xi^-)$ and $s(\xi) > t(\xi)$ where ξ^- is the greatest element of F less than ξ .⁹*

The following notation will be convenient.

Definition. *If $\alpha < \beta < \omega_1$, let $\text{Osc}(\alpha, \beta)$ denote*

$$\text{Osc}(e_\alpha, e_\beta; L(\alpha, \beta))$$

and $\text{osc}(\alpha, \beta)$ denote the cardinality of $\text{Osc}(\alpha, \beta)$.

Lemma 4.1. *For every uncountable pairwise disjoint $\mathcal{A} \subseteq [\omega_1]^k$ and $\mathcal{B} \subseteq [\omega_1]^l$ and all w in $C(2^\omega, \omega)$, there are b_m ($m < \omega$) in $\mathcal{B}$ such that for all n there is an a in $\mathcal{A}$ and $\{\xi_i : i < n\}$ such that for all $m \leq n$, $i < k$, $j < l$ the following hold:*

⁸A subset of ω_1 is *club* if it is closed and unbounded. A subset of ω_1 is *stationary* if it intersects every club.

⁹Actually we will later see that we are really interested in counting oscillations between the relations $=$ and $\neq$ rather than the more conventional $<$ and $>$. The above definition is a compromise between the two.

- (1) $a < b_m$.
- (2) $\text{Osc}(a(i), b_m(j))$ is the disjoint union of $\text{Osc}(a(i), b_0(j))$ and $\{\xi_{m'} : m' < m\}$.
- (3) $\mu(a(i), b_0(j))$ is $\mu(a(i), b_m(j))$ restricted to $L(a(i), b_0(j))$.
- (4) $\mu(a(i), b_m(j))(\xi_{m'}) = w$ whenever $m' < m$.

Remark. While the generality of this theorem will be needed to derive some of the results in the paper, many of them can get by with weaker formulations. For instance, the analysis of osc only requires the theorem for $k = l = 1$ and only conclusions (1) and (2). This is sufficient to prove Theorem 1.10. The later conditions are needed for the properties of the stronger coloring o presented in Section 5.

Lemma 4.1 will be derived from the following theorem.

Lemma 4.2. *Let $\mathcal{A} \subseteq [\omega_1]^k$ and $\mathcal{B} \subseteq [\omega_1]^l$ be uncountable and pairwise disjoint. There is a closed and unbounded set of $\delta < \omega_1$ such that if a is in $\mathcal{A} \setminus \delta$,¹⁰ b is in $\mathcal{B} \setminus \delta$, and R is in $\{=, >\}$, then there are a^+ in $\mathcal{A} \setminus \delta$, and b^+ in $\mathcal{B} \setminus \delta$ such that for all $i < k$ and $j < l$:*

- (1) *The maximum of $L(\delta, b(j))$ is less than both $\Delta(e_{a(i)}, e_{a^+(i)})$ and $\Delta(e_{b(j)}, e_{b^+(j)})$.*
- (2) *There is a non-empty L^+ which does not depend on j such that $L(\delta, b(j)) < L^+$ and $L(\delta, b^+(j)) = L(\delta, b(j)) \cup L^+$.*
- (3) *If ξ is in L^+ , then $e_{a^+(i)}(\xi) R e_{b^+(j)}(\xi)$.*
- (4) *$\mu(\delta, b(j))$ is $\mu(\delta, b^+(j))$ restricted to $L(\delta, b(j))$.*
- (5) *$\mu(\delta, b^+(j))(\min L^+) = w_\delta$.*

Proof. Let M be an elementary submodel of $H(\aleph_2)$ which contains everything relevant and let $\delta = M \cap \omega_1$. By Fact 6, it suffices to show that δ satisfies the conclusion of the theorem.

First suppose that R is $=$. Applying Fact 5, find a $\gamma_0 < \delta$ satisfying the following conditions:

- (1) If $j < l$, then $L(\delta, b_m(j)) < \gamma_0$.
- (2) If $i < k$, $j < l$, and $\gamma_0 < \xi < \delta$, then $e_{a(i)}(\xi) = e_{b(j)}(\xi)$.

Applying Facts 2 and 4, pick a $\gamma < \delta$ such that if $\gamma < \xi < \delta$, then $\gamma_0 < L(\xi, \delta)$. Consider the set $D \subseteq \omega_1$ consisting of all δ^+ such that for some a^+ in $\mathcal{A} \setminus \delta^+$ and b^+ in $\mathcal{B} \setminus \delta^+$ the following conditions are satisfied:

- (1) $e_{a^+(i)} \upharpoonright \gamma_0 = e_{a(i)} \upharpoonright \gamma_0$ and $e_{b^+(j)} \upharpoonright \gamma_0 = e_{b(j)} \upharpoonright \gamma_0$ for all $i < k$ and $j < l$.

¹⁰It will be convenient to let $\mathcal{A} \setminus \delta$ denote the set of all a in $\mathcal{A}$ such that $\delta \leq a$ whenever $\mathcal{A}$ is a collection of finite subsets of ω_1 and $\delta < \omega_1$.

- (2) If $\gamma < \xi < \delta^+$, then $\gamma_0 < L(\xi, \delta^+)$ and $\mu(\xi, \delta^+)(\min L(\xi, \delta^+)) = w_{\delta^+} = w_\delta$.
- (3) If $\gamma_0 < \xi < \delta^+$, $i < k$, and $j < l$, then $e_{a^+(i)}(\xi) = e_{b^+(j)}(\xi)$.
- (4) $\mu(\delta^+, b^+(j)) = \mu(\delta, b(j))$.

Observe that for all $\beta \geq \gamma_0$, $e_\beta \upharpoonright \gamma_0$ is in M since by Fact 5 it differs from e_{γ_0} on a finite set. Hence D is definable from the parameters γ_0 , γ , $e_{a(i)} \upharpoonright \gamma_0$, etc. which are all elements of M . Therefore D is in M by Fact 9. Since D has δ as a member, it is uncountable by Fact 7. Hence there is a $\delta^+ > \delta$ in D . Let $a^+ \in \mathcal{A} \setminus \delta^+$ and $b^+ \in \mathcal{B} \setminus \delta^+$ witness that δ^+ is in D .

Now let $i < k$ and $j < l$ be arbitrary. First observe that

$$\gamma_0 \leq \Delta(e_{a(i)}, e_{a^+(i)}),$$

$$\gamma_0 \leq \Delta(e_{b(j)}, e_{b^+(j)}).$$

Put $L^+ = L(\delta, \delta^+)$. Notice that $L(\delta, b(j)) < L^+$ and hence

$$L(\delta, b^+(j)) = L(\delta^+, b^+(j)) \cup L^+$$

holds by Fact 1. Since $\delta < \delta^+$, L^+ is non-empty. If ξ is in L^+ , then $\gamma_0 < \xi < \delta^+$ and so

$$e_{a^+(i)}(\xi) R e_{b^+(j)}(\xi).$$

Now suppose that R is $>$. Put E equal to the set of all limit $\nu < \omega_1$ such that for all a_0 in $\mathcal{A} \setminus \nu$, all $\nu_0 < \nu$, $\varepsilon < \omega_1$, $n < \omega$, and finite $L^+ \subseteq \omega_1 \setminus \nu$ there is an a_1 in $\mathcal{A} \setminus \varepsilon$ with

$$\nu_0 \leq \Delta(e_{a_0(i)}, e_{a_1(i)}),$$

$$e_{a_1(i)}(\xi) > n$$

whenever $i < k$ and ξ is in L^+ . By Fact 9, E is an element of M .

Claim 4.3. δ is in E . In particular, E is uncountable by Fact 7.

Proof. Let a_0 , ν_0 , ε , n , and L^+ be given as in the definition of E for $\nu = \delta$. By Fact 5 we may assume without loss of generality that ν_0 is an upper bound for all $\xi < \delta$ such that $e_{a_0(i)}(\xi) \leq n$ for some $i < k$. Now, applying elementarity of M , there is a δ^+ above ε , δ , and $\max L^+$ and an a_1 in $\mathcal{A} \setminus \delta^+$ such that the following conditions are satisfied:

- (1) For all $i < k$, $e_{a_0(i)} \upharpoonright \nu_0 = e_{a_1(i)} \upharpoonright \nu_0$.
- (2) If $\nu_0 < \xi < \delta^+$ and $i < k$, then $e_{a_1(i)}(\xi) > n$.

Since $L^+ \subseteq \delta^+ \setminus \delta$, this completes the proof of the claim. $\square$

Applying elementarity of M and uncountability of E , find an element γ_0 of E such that $L(\delta, b(j)) < \gamma_0 < \delta$ for all $j < l$. Applying Fact 2, find a $\gamma < \delta$ such that if $\gamma < \xi < \delta$, then $\gamma_0 < L(\xi, \delta)$. Again using

elementarity of M , select a limit $\delta^+ > \delta$ and a b^+ in $\mathcal{B} \setminus \delta^+$ so that the following conditions are satisfied:

- (1) $e_{b^+(j)} \upharpoonright \gamma_0 = e_{b(j)} \upharpoonright \gamma_0$ for all $j < l$.
- (2) If $\gamma < \xi < \delta^+$, then $\gamma_0 < L(\xi, \delta^+)$ and $\mu(\xi, \delta^+)(\min L(\xi, \delta^+)) = w_{\delta^+} = w_\delta$.
- (3) $\mu(\delta^+, b^+(j)) = \mu(\delta, b(j))$.

Put $L^+ = L(\delta, \delta^+)$. Applying the definition of E , find an a^+ in $\mathcal{A} \setminus \delta$ such that for all $i < k$, $j < l$, and ξ in L^+

$$L(\delta, b(j)) < \Delta(e_{a(i)}, e_{a^+(i)})$$

$$e_{a^+(i)}(\xi) > e_{b^+(j)}(\xi).$$

The rest of the verification is as in the previous case. This completes the proof of Lemma 4.2. $\square$

Now we are ready to prove Lemma 4.1

Proof. Let $\mathcal{A}$ and $\mathcal{B}$ be given and select a countable elementary submodel M of $H(\aleph_2)$ containing everything relevant such that $w_\delta = w$ where $\delta = M \cap \omega_1$. Since M contains $\mathcal{A}$ and $\mathcal{B}$, the closed and unbounded set provided by Lemma 4.2 is in M and therefore δ is in this closed and unbounded set. Using Lemma 4.2, it is easy to select a_m ($m < \omega$) in $\mathcal{A} \setminus \delta$ and b_m ($m < \omega$) in $\mathcal{B} \setminus \delta$ so that for all $m < \omega$ the following conditions are satisfied whenever $i < k$ and $j < l$:

- (1) $L(\delta, b_{m+1}(j))$ is a proper initial part of $L(\delta, b_m(j))$.
- (2) The minimum of $L(\delta, b_{m+1}(j)) \setminus L(\delta, b_m(j))$ is an ordinal ξ_m which does not depend on j .
- (3) $\text{Osc}(a_{m+1}(i), b_{m+1}(j); L(\delta, b_{m+1}(j)))$ is formed by adding ξ_m to $\text{Osc}(a_m(i), b_m(j); L(\delta, b_m(j)))$.
- (4) If $m' < m$, then $\xi_{m'}$ is less than both $\Delta(e_{a_m(i)}, e_{a_{m+1}(i)})$ and $\Delta(e_{b_m(j)}, e_{b_{m+1}(j)})$.
- (5) $e_{a_m(i)}(\max L(\delta, b_m(j))) > e_{b_m(j)}(\max L(\delta, b_m(j)))$.
- (6) $\mu(\delta, b_0(j))$ is the restriction of $\mu(\delta, b_m(j))$ to $L(\delta, b_0(j))$.
- (7) If $m' < m$, then $\mu(\delta, b_m(j))(\xi_{m'}) = w$.

Now let n be given. Pick a $\gamma_0 < \delta$ which is an upper bound for each $L(\delta, b_n(j))$ for $j < l$ and all $\xi < \delta$ such that for some $m, m' \leq n$ and $j < l$

$$e_{b_m(j)}(\xi) \neq e_{b_{m'}(j)}(\xi)$$

(the later set is finite by Fact 5). Using elementarity of M and Fact 2, select an a in $\mathcal{A}$ such that $a < \delta$ and for all $i < k$ and $j < l$

$$L(\delta, b_n(j)) < \Delta(e_{a(i)}, e_{a_n(i)}),$$

$$\gamma_0 < L(a(i), \delta).$$

Now let $i < k$, $j < l$, and $m \leq n$ be fixed. It follows from Fact 1 that

$$\mu(a(i), b_m(j)) = \mu(a(i), \delta) \cup \mu(\delta, b_m(j)).$$

Also,

$$\mu(a(i), b_m(j))(\xi_{m'}) = \mu(\delta, b_m(j))(\xi_{m'}) = w.$$

Finally, $e_{b_m(j)} \upharpoonright L(a(i), \delta)$ does not depend on m and therefore

$$\text{Osc}(a(i), b_0(j); L(a(i), \delta)) = \text{Osc}(a(i), b_m(j); L(a(i), \delta)).$$

Hence, by item 5,

$$\text{Osc}(a(i), b_m(j)) = \text{Osc}(a(i), b_0(j)) \cup \{\xi_{m'} : m' < m\}.$$

This completes the proof of Lemma 4.1. $\square$

5. A NEGATIVE PARTITION RELATION AND THE COLORING o

We will begin this section with an important consequence of Lemma 4.1. First I will introduce some notation.

Definition. The function $*$: $\omega \rightarrow \omega$ is defined by letting $*(0) = 0$ and if $m > 0$, then $*(m) = n$ if the n^{th} prime is the least prime which does not divide m . If f is a function taking values in ω , then f^* will denote the composition of f followed by $*$.

The notation for $*$ is chosen to mimic its usage in [33]. In particular, the following is easily verified.

Fact 10. If $X \subseteq \omega$ contains arbitrarily long intervals of integers, then the image of X under $*$ is ω .

Hence Lemma 4.1 immediately yields the following result.

Theorem 5.1. If $A, B \subseteq \omega_1$ are uncountable and $n < \omega$, then there are α in A and β in B with $\alpha < \beta$ and $\text{osc}^*(\alpha, \beta) = n$.

In other words, osc^* witnesses $\omega_1 \not\rightarrow [\omega_1; \omega_1]_{\omega}^2$. If $\iota_\beta : \beta \rightarrow \omega$ is an injection for each $\beta < \omega_1$, then for $\alpha < \beta$ we can put $f(\alpha, \beta) = \xi$ if $\text{osc}^*(\alpha, \beta) = \iota_\beta(\xi)$ and $f(\alpha, \beta) = 0$ if no such ξ exists. It is easily checked this witnesses the negative partition relation in the following theorem.

Theorem 5.2. $\omega_1 \not\rightarrow [\omega_1; \omega_1]_{\omega_1}^2$.

Two variations of $\omega_1 \rightarrow (\omega_1; \omega_1)_2^2$ which are closely related to (S) and (L) respectively are:

$$\omega_1 \rightarrow (\omega_1, (\omega_1; [\omega_1]^{<\aleph_0}))^2,$$

$$\omega_1 \rightarrow (\omega_1, ([\omega_1]^{<\aleph_0}; \omega_1))^2.$$

The statement $\omega_1 \rightarrow (\omega_1, (\omega_1; [\omega_1]^{<\aleph_0}))^2$ asserts that if $c : [\omega_1]^2 \rightarrow 2$ then either

- (1) there is an uncountable $X \subseteq \omega_1$ such that c is constantly 0 on $[X]^2$ or
- (2) there are uncountable $A \subseteq \omega_1$ and uncountable pairwise disjoint $\mathcal{B} \subseteq [\omega_1]^{<\aleph_0}$ such that for all α in A and b in $\mathcal{B}$ with $\alpha < b$, there is a β in b with $c(\alpha, \beta) = 1$.

The statement $\omega_1 \rightarrow (\omega_1, ([\omega_1]^{<\aleph_0}; \omega_1))^2$ is similarly defined by replacing $A \subseteq \omega_1$ by a pairwise disjoint $\mathcal{A} \subseteq [\omega_1]^{<\aleph_0}$ and $\mathcal{B}$ by $B \subseteq \omega_1$. These imply (S) and (L) respectively (see [27]) and are each consequences of $\omega_1 \rightarrow (\omega_1; \omega_1)_2^{11}$.

In [27], Todorćević showed that $\omega_1 \rightarrow (\omega_1, (\omega_1; [\omega_1]^{<\aleph_0}))^2$ was relatively consistent with ZFC and in fact follows from PFA. We will now consider a map $o : [\omega_1]^2 \rightarrow \omega$ such that o^* witnesses a strong failure of $\omega_1 \rightarrow (\omega_1, ([\omega_1]^{<\aleph_0}; \omega_1))^2$.

Definition. Let $o(\alpha, \beta)$ denote

$$\sum_{q \neq 0} (|\mu(\alpha, \beta; \alpha)^{-1}(q)| \bmod q).$$

Theorem 5.3. Let $\mathcal{A} \subseteq [\omega_1]^k$ and $\mathcal{B} \subseteq [\omega_1]^l$ be uncountable and pairwise disjoint families. For every $\chi : k \rightarrow 2$ and $\pi : k \rightarrow l$ there are a in $\mathcal{A}$ and b in $\mathcal{B}$ such that $a < b$ and for all $i < k$,

$$o^*(a(i), b(\pi(i))) = \chi(i).$$

Remark. It is possible to prove this result if 2 is replaced by any positive integer. We will only need this formulation and, alas, we are running out of good letters for integer variables.

Proof. Let $\mathcal{A}$, $\mathcal{B}$, χ , and π be given as in the statement of the theorem. Select distinct primes q_i ($i < k$) which are each larger than 8. By refining $\mathcal{A}$ if necessary, we may assume that there is a w in $C(2^\omega, \omega)$ such that if a is in $\mathcal{A}$, then $w(a(i)) = q_i$ for all $i < k$. Applying Lemma 4.1, find $a \in \mathcal{A}$, $b_m \in \mathcal{B}$, ξ_m for each m less than $n = 6 \cdot \prod_{i < k} q_i$ such that the following conditions are satisfied for all $m < n$, $i < k$, and $j < l$:

- (1) $a < b_m$.

¹¹Of course this latter statement is trivial in light of Theorem 1.10 but this implication was already observed by the 1980's.

- (2) $\text{Osc}(a(i), b_m(j))$ is the disjoint union of the sets $\text{Osc}(a(i), b_0(j))$ and $\{\xi_{m'} : m' < m\}$.
- (3) $\mu(a(i), b_0(j))$ is $\mu(a(i), b_m(j))$ restricted to $L(a(i), b_0(j))$.
- (4) $\mu(a(i), b_m(j))(\xi_{m'}) = w$ whenever $m' < m$.

Put

$$r_i = \left(\sum_{s \neq 0, q_i} |\mu(a(i), b_0(\pi(i)); a(i))^{-1}(s)| \bmod s \right) \bmod 6.$$

Use the Chinese Remainder Theorem to find an x which satisfies the following equations for each $i < k$:

$$x + |\mu(a(i), b_0(\pi(i)); a(i))^{-1}(q_i)| \equiv 6 - r_i + 2^{\chi(i)} \pmod{q_i}.$$

Notice that the right hand side of these equations are always at least 0 and strictly less than q_i since $8 < q_i$. Let $b = b_x$.

We now need to show that for each $i < k$, the least prime which does not divide $o(a(i), b(\pi(i)))$ is 2 if $\chi(i) = 0$ and 3 otherwise. Let $i < k$ be fixed. Notice that, for some y

$$\begin{aligned} o(a(i), b(\pi(i))) &= \left(x + \mu(a(i), b_0(\pi(i)); a(i))^{-1}(q_i) \right) \bmod q_i + r_i + 6y \\ &= (6 - r_i + 2^{\chi(i)}) + r_i + 6y \\ &= 2^{\chi(i)} + 6(y + 1). \end{aligned}$$

Since 3 divides $6(y + 1)$ but not $2^{\chi(i)}$, 3 does not divide $o(a(i), b(\pi(i)))$. On the other hand, 2 divides $o(a(i), b(\pi(i)))$ if and only if $\chi(i) = 1$. It follows that $o^*(a(i), b(\pi(i))) = \chi(i)$. $\square$

Notice that we can think of the above result in the following way. We say that χ is a *pattern* if for some positive integers k and l and some subset $D \subseteq k \times l$, χ is a function from D into ω . A coloring $c : [\omega_1]^2 \rightarrow \omega$ is said to always realize a pattern χ if whenever $\mathcal{A} \subseteq [\omega_1]^k$ and $\mathcal{B} \subseteq [\omega_1]^l$ are pairwise disjoint and uncountable, there are a in $\mathcal{A}$ and b in $\mathcal{B}$ such that $a < b$ and

$$c(a(i), b(j)) = \chi(i, j)$$

whenever (i, j) is in D .

Theorem 5.3 just says that the coloring o^* always realizes every binary pattern in which D defines a function. On the other hand, Todorćević has shown that PFA implies $\omega_1 \rightarrow (\omega_1, (\omega_1; [\omega_1]^{<\aleph_0}))^2$ which implies that for every coloring $c : [\omega_1]^2 \rightarrow 2$, there is a constant pattern on $1 \times l$ which is not always realized by c . This suggests the following question.

Question 5.4. (PFA) *If $c : [\omega_1]^2 \rightarrow 2$, is there a pattern on 1×2 which is not always realized?*

The proof of Theorem 7.9 below shows that o cannot provide a counterexample to this question.

6. A FAMILY OF BINARY RELATIONS ON ω_1

In this section we will consider the family of binary relations on ω_1 with the Tukey order. Recall that the definition of the Tukey order $\leq$ on binary relations R and S .

Definition. *If R and S are binary relations, then we write $R \leq S$ iff there are functions $f : \text{dom}(R) \rightarrow \text{dom}(S)$ and $g : \text{ran}(S) \rightarrow \text{ran}(R)$ such that*

$$f(x) S y \text{ implies } x R g(y).$$

This order was first considered by J.W. Tukey in the class of transitive relations [34]. It make sense, however, to consider this relation on the more general class of binary relations (see [35]).

Our focus will be on binary relations on ω_1 . Two fundamental examples are the well order ω_1 and the family $[\omega_1]^{<\aleph_0}$ of finite subsets of ω_1 ordered by inclusion. We will also need a few standard operations on relations. If R and S are relations, then $R \oplus S$ is the relation which is the disjoint union of R and S . If m is a cardinal and R is a relation, then we will let $m \cdot R$ will denote the direct sum of m copies of R . If R and S are two binary relations, then $R \wedge S$ and $R \vee S$ are the relations with domains $\text{dom}(R) \times \text{dom}(S)$ and ranges $\text{ran}(R) \times \text{ran}(S)$ such that

$$(a, b) R \wedge S (c, d) \text{ iff } a R c \text{ or } b S d,$$

$$(a, b) R \vee S (c, d) \text{ iff } a R c \text{ and } b S d.$$

The join $R \vee S$ is often denoted $R \times S$. It is easily verified that $\wedge$ and $\vee$ give lower and upper bounds respectively.¹²

It was observed that the binary relations on ω_1 which one knew how to construct in ZFC were either below $\aleph_0 \cdot \omega_1$ or above $[\omega_1]^{<\aleph_0}$ in the Tukey order.

Example 6.1. *Define $\alpha R \beta$ iff $\alpha < \beta$ and $z_\alpha < z_\beta$.*

The relation R is essentially Sierpiński's partition which witnesses $\omega_1 \not\leq (\omega_1)_2^2$. In this case $R \leq \aleph_0 \cdot \omega_1$ by the following Tukey maps:

$$f(\alpha) = (\min\{n : z_\alpha < q_n\}, \alpha)$$

$$g(\alpha, n) = \min\{\beta : \alpha < \beta \wedge q_n < z_\beta\}$$

¹²In general these need not be optimal — they are in the class of directed relations.

where $\{q_n\}_{n < \omega}$ is an enumeration of a countable order-dense subset of 2^ω . On the other hand the following example shows that with an additional hypothesis such as $\diamond$, one can construct more complex relations.

Example 6.2. *Let R be a tree order on a set T . If $R \leq \aleph_0 \cdot \omega_1$, then T is the union of countably many R -chains. If $[\omega_1]^{<\aleph_0} \leq R$, then T contains an uncountable R -antichain. In particular, if (T, R) is a Suslin tree, then $R \not\leq \aleph_0 \cdot \omega_1$ and $[\omega_1]^{<\aleph_0} \not\leq R$.*

These observations led to the following conjecture.

Conjecture 6.3. (PFA) *If R is a binary relation, then either $R \leq \aleph_0 \cdot \omega_1$ or else $[\omega_1]^{<\aleph_0} \leq R$.*

This can be considered a basis conjecture in the following sense. Let $\mathcal{R}$ denote the class of all binary relations R on ω_1 such that R is not reducible to $\aleph_0 \cdot \omega_1$. Then the above conjecture is just the assertion that $\mathcal{R}$ has a single element basis consisting of $[\omega_1]^{<\aleph_0}$. This conjecture was given further plausibility by the following theorem which implies that it is true for transitive relations.

Theorem 6.4. [32] (PFA) *Every transitive relation on ω_1 is Tukey equivalent to one of the following for some non negative integers n_i ($i < 5$):*

- (1) $n_0 \cdot 1 \oplus n_1 \cdot \omega \oplus n_2 \cdot \omega_1 \oplus n_3 \cdot \omega \times \omega_1 \oplus n_4 \cdot [\omega_1]^{<\aleph_0}$.
- (2) $\aleph_0 \cdot 1 \oplus n_2 \cdot \omega_1 \oplus n_3 \cdot \omega \times \omega_1 \oplus n_4 \cdot [\omega_1]^{<\aleph_0}$.
- (3) $\aleph_0 \cdot \omega_1 \oplus n_4 \cdot [\omega_1]^{<\aleph_0}$.
- (4) $\aleph_0 \cdot [\omega_1]^{<\aleph_0}$.
- (5) $=$.

We will now see, however, that Conjecture 6.3 is provably false.

Definition. Let $c(\alpha, \beta)$ denote $o(\alpha, \beta) \bmod 2$ if $\alpha < \beta$.

Definition. Let R denote the relation with domain and range ω_1 defined by letting $\alpha R \beta$ iff $\alpha = \beta$ or $\alpha < \beta$ and $c(\alpha, \beta) = 1$. If X is an uncountable subset of ω_1 , let R_X denote the relation with domain X and range ω_1 which is the restriction of R to $X \times \omega_1$.

Trivially, $\omega_1 \leq R_X$ for every uncountable X and if $X \subseteq Y$, then $R_X \leq R_Y$. The following shows that none of these relations are above $[\omega_1]^{<\aleph_0}$ in the Tukey order.

Theorem 6.5. $[\omega_1]^{<\aleph_0} \not\leq R$.

Proof. Suppose that $f : [\omega_1]^{<\aleph_0} \rightarrow \omega_1$ and $g : \omega_1 \rightarrow [\omega_1]^{<\aleph_0}$ are functions. It suffices to show that they are not Tukey reductions.

Applying the pressing down lemma, find a stationary $B \subseteq \omega_1$ and a finite $x_0 \subseteq \omega_1$ such that for all β in B

$$g(\beta) \cap \beta = x_0.$$

Let

$$\begin{aligned}\mathcal{X} &= \{x \in [\omega_1]^{<\aleph_0} : x \not\subseteq x_0\}, \\ A &= \{f(x) : x \in \mathcal{X}\}.\end{aligned}$$

By refining B if necessary, we may assume that if α is in A and β is in B with $\alpha < \beta$, then there is an x in $\mathcal{X}$ such that $f(x) = \alpha$ and $x \subseteq \beta$.

If A is countable, then we can find a α in A such that

$$\{x \in \mathcal{X} : f(x) = \alpha\}$$

is uncountable. It is then easy to find an x in $\mathcal{X}$ such that x is not contained in $g(\alpha)$ but $f(x) = \alpha$ and hence $f(x) R \beta$, witnessing that f, g are not Tukey maps.

If A is uncountable, apply Theorem 5.3 to obtain an α in A and a β in B such that $\alpha < \beta$ and $c(\alpha, \beta) = 1$. Now pick an x in $\mathcal{X}$ such that $f(x) = \alpha$ and $x \subseteq \beta$ and observe that x is not contained in $g(\beta)$ and yet $f(x) R \beta$. $\square$

Theorem 6.6. *For all finite sets F of uncountable subsets of ω_1 ,*

$$\bigwedge_{X \in F} R_X \not\leq \aleph_0 \cdot \omega_1.$$

Proof. By replacing F with a disjoint refinement if necessary, we may assume without loss of generality that F consists of pairwise disjoint sets. Suppose for contradiction that

$$f : \prod_{X \in F} X \rightarrow \omega \times \omega_1$$

$$g : \omega \times \omega_1 \rightarrow \prod_{X \in F} \omega_1$$

are Tukey maps. Find uncountable pairwise disjoint $\mathcal{A}, \mathcal{B} \subseteq [\omega_1]^{|F|}$ and an n such that:

- (1) For each element a of $\mathcal{A} \cup \mathcal{B}$, $a \cap X$ is a singleton for all X in F .
- (2) $f \upharpoonright \mathcal{A}$ is an injection into $\{n\} \times \omega_1$.
- (3) $g^{-1}\mathcal{B} \subseteq \{n\} \times \omega_1$.

Applying Theorem 5.3, it is possible to find an a in $\mathcal{A}$ and an x in $g^{-1}\mathcal{B}$ such that $a < g(x)$, $f(a) < x$, and for all X in F

$$c(a \cap X, g(x) \cap X) = 0$$

(where we are identifying the singletons $a \cap X$ and $g(x) \cap X$ with their element). It follows immediately that $f(a) \leq x$ while a is not $\bigwedge_{X \in F} R_X$ -related to $g(x)$, a contradiction. $\square$

The next theorem show that these relations are typically incomparable.

Theorem 6.7. *If $X \setminus Y$ is uncountable, then $R_X \not\leq R_Y$.*

Proof. Suppose for contradiction that $f : X \rightarrow Y$ and $g : \omega_1 \rightarrow \omega_1$ are Tukey maps. To obtain a contradiction, it suffices to find α in X and a β in ω_1 such that $f(\alpha) < \beta$, $\alpha < g(\beta)$, $c(f(\alpha), \beta) = 1$, and $c(\alpha, g(\beta)) = 0$. Since f and g are Tukey reductions it is possible to find an uncountable $X_0 \subseteq X \setminus Y$ and uncountable $Z \subseteq \omega_1$ such that $f \upharpoonright X_0$ and $g \upharpoonright Z$ are injections into sets disjoint from X_0 and Z , respectively, and the inequalities $\alpha < f(\alpha)$, $\beta = f(\beta)$, $\beta < f(\beta)$ are uniformly true or false as α ranges over X_0 and β ranges over Z . Put

$$\mathcal{A} = \{\{\alpha, f(\alpha)\} : \alpha \in X_0\}$$

$$\mathcal{B} = \{\{\beta, g(\beta)\} : \beta \in Z\}$$

and notice that these families are uncountable and consist of pairwise disjoint sets of uniform cardinality which, in the case of $\mathcal{A}$, is two. By our uniformity assumption on the relations $<$ and $=$, it is possible to find i_0 and j_0 such that if a is in $\mathcal{A}$, then $a(i_0)$ is in X_0 and if b is in $\mathcal{B}$, then $b(j_0)$ is in Z . Put $\pi(i_0) = j_0$ and $\pi(1 - i_0) = 1 - j_0$ if $\mathcal{B}$ consists of doubletons and $\pi(1 - i_0) = j_0$ otherwise. Define $\chi(i_0) = 0$ and $\chi(1 - i_0) = 1$. Applying Theorem 5.3 to $\mathcal{A}$, $\mathcal{B}$, π , and χ , there are a in $\mathcal{A}$ and b in $\mathcal{B}$ such that

$$a < b$$

$$c(a(i), b(\pi(i))) = \chi(i).$$

Translating the outcome, it is easily checked that $\alpha = a(i_0)$ and $\beta = b(j_0)$ are as desired. $\square$

The following remain unclear.

Question 6.8. *For which families $\mathcal{F} \subseteq [\omega_1]^{\aleph_1}$ is there an S such that $S \not\leq \aleph_0 \cdot \omega_1$ and $S \leq R_X$ for all X in $\mathcal{F}$.*

Question 6.9. *Is the collection of all S with $S \not\leq \aleph_0 \cdot \omega_1$ downwards-directed in the Tukey order?*

7. AN L SPACE AND THE NON-EXISTENCE OF A SMALL BASIS FOR THE REGULAR HAUSDORFF SPACES.

In this section I will give an example of an *L space* — a hereditarily Lindelöf, not hereditarily separable space.¹³ The question of the existence of such spaces was first asked explicitly in [9], though arguably this question can be traced to Suslin's [25] where he posed his famous hypothesis. For instance, an immediate consequence of Kurepa's [16] is that a Suslin line is an example of an L space. From the 1960's until the 1980's, there was a concerted effort to understand both L spaces and their dual the S space. I refer the reader to Juhasz's [12] and Rudin's [21] as well as Roitman's more recent [20] for more discussion on these developments. I will mention a few. First, Roitman and Zenor showed that there was a relationship between the existence of certain S and L spaces.

Theorem 7.1. [19] [37] *There is a strong S space iff there is a strong L space.*¹⁴

Hence the difference in the existence of S and L spaces lies in the properties of their finite powers. This gives some explanation as to why the existence of S and L spaces seem to be such similar hypotheses at first.

There are a number of results under $\text{MA}_{\aleph_1}$ which limit the existence of S and L spaces.

Theorem 7.2. [13] $(\text{MA}_{\aleph_1})$ *There are no strong S or L spaces.*

Theorem 7.3. [26] $(\text{MA}_{\aleph_1})$ *There are no first countable L spaces.*

Theorem 7.4. [6] $(\text{MA}_{\aleph_1})$ *If K is a compact space which contains an L space, then K maps continuously onto $[0, 1]^{\aleph_1}$.*

Finally, Todorćević proved that S spaces do not exist assuming PFA.

Theorem 7.5. [27] (PFA) *If a space is hereditarily separable, then it is hereditarily Lindelöf.*

Until now, however, it remained unclear whether Todorćević's methods could be used to prove that PFA implies there are no L spaces. I will now show that this is not the case — that there is an L space

¹³According to Juhasz, M.E. Rudin coined the phrase “S space” but to mean an hereditarily separable space. Juhasz then introduced the phrase “L space” and gave the current definitions to S and L spaces as those which distinguish hereditary separability from the hereditary Lindelöf property. Also, it was shown in [9] that examples exist without the assumption that the spaces are regular and Hausdorff.

¹⁴Here a space is a strong S (L) space iff all of its finite powers are S (L) spaces.

which can be constructed without appealing to additional axioms of set theory. For each $\alpha < \omega_1$, put

$$W_\alpha = \{\alpha\} \cup \{\beta > \alpha : c(\alpha, \beta) = 1\}.$$

If $X \subseteq \omega_1$, then let $\tau[X]$ be the topology on X generated by declaring $W_\xi \cap X$ to be clopen for all ξ in X . If X is uncountable, then clearly $\tau[X]$ is non-separable. The following theorem shows that $\tau[X]$ is always Fréchet.

Theorem 7.6. *If X is a set and $\mathcal{F}$ is a point countable point separating family of subsets of X , then the topology on X defined by declaring elements of $\mathcal{F}$ to be clopen is countably tight and has every countable subspace metrizable. In particular the topology is Fréchet.*

Remark. I attribute this result to Zoltan Balogh. He once told me that “If there is an L space, then there is a countably tight one.” While I never saw his proof, I think it is reasonable to assume this he may have proceeded along these lines.

Proof. That countable subspaces are metrizable follows from the assumptions and the well known fact that a (regular Hausdorff) second countable spaces are metrizable. To see that the space is countably tight, let $A \subseteq X$ have an accumulation point x in X . Let M be a countable elementary submodel of $H(\theta)$ for θ regular and large enough so that X , $\mathcal{F}$, x , and A are all in M . It suffices to show that x is an accumulation point of $A \cap M$.

Suppose this is not the case and let U_i ($i < k$) and V_i ($i < l$) be elements of $\mathcal{F}$ such that

$$W = \bigcap_{i < k} U_i \setminus \bigcup_{j < l} V_j$$

contains x and is disjoint from $A \cap M$. The important observation is this: if V is in $\mathcal{F}$, then $V \cap M$ is non-empty iff V is in M . This is because $\{y \in V : V \in \mathcal{F}\}$ is countable for all y in X and therefore a subset of M , 9) whenever y is in M (by Facts 7. Hence each U_i must be in M since x is in U_i and x is in M . Moreover, since we are only interested in having W be disjoint from $A \cap M$, we may assume without loss of generality that each V_j is in M . But then W must be in M and, by elementarity of M , there must be an element of $W \cap A$ which is in M , a contradiction. $\square$

It is straightforward to use Theorem 5.3 to show that every $\tau[X]$ is hereditarily Lindelöf; I will prove a result of independent interest and then derive this as a consequence.

Theorem 7.7. *If $X, Y \subseteq \omega_1$ have countable intersection, then there is no continuous injection from any uncountable subspace of $(X, \tau[X])$ into $(Y, \tau[Y])$.*

Proof. Without loss of generality X and Y are actually disjoint. Suppose for contradiction that such an injection $f : X_0 \rightarrow Y$ does exist for some uncountable $X_0 \subseteq X$. Let $\mathcal{B}$ be the set of all $\{\beta, f(\beta)\}$ such that β is in X_0 . For simplicity we will assume that $\beta < f(\beta)$ for all β ; a similar argument is given if $f(\beta) < \beta$ for uncountably many β . For each β in X_0 , pick a basic open set U_β which contains β such that $f^{-1}W_{f(\beta)}$ contains U_β . By refining X_0 if necessary, we may assume that for some k and $\chi_0 : k \rightarrow 2$ there is a Δ -system $\{F_\beta : \beta \in X_0\}$ with root F of $|F| + k$ -element subsets of X such that:

- (1) If $F_\alpha < \beta$, then β is in U_α iff for all $i < |F| + k$, $c(F_\alpha(i), \beta) = \chi_0(i)$.
- (2) $l = |F_\alpha \cap f(\alpha)|$ does not depend on α .
- (3) $\max F$ is less than both α and $\min(F_\alpha \setminus F)$.

Let $\mathcal{A}$ be the collection of all sets of the form $F_\alpha \setminus F \cup \{f(\alpha)\}$. Notice that since X is disjoint from Y , $f(\alpha)$ is not in F_α for any α in X_0 . Define $\chi : k + 1 \rightarrow 2$ by $\chi(i) = \chi_0(|F| + i)$ if $i < l$, $\chi(l) = 0$, and $\chi(i) = \chi_0(|F| + i - 1)$ if $i > l$. Define $\pi : k + 1 \rightarrow 2$ by $\pi(i) = 0$ if $i \neq l$ and $\pi(l) = 1$. Applying Theorem 5.3, there are $a \in \mathcal{A}$ and $b \in \mathcal{B}$ such that $a < b$ and for all $i < k + 1$

$$c(a(i), b(\pi(i))) = \chi(i).$$

Let α, β in X_0 be such that

$$\begin{aligned} a &= (F_\alpha \setminus F) \cup \{f(\alpha)\} \\ b &= \{\beta, f(\beta)\}. \end{aligned}$$

To derive a contradiction, it suffices to show that β is in U_α but that $f(\beta)$ is not in $W_{f(\alpha)}$. The latter is just a reformulation of

$$c(a(l), b(\pi(l))) = c(f(\alpha), f(\beta)) = \chi(l) = 0.$$

For the former, we need to show that for all $i < |F| + k$,

$$c(F_\alpha(i), \beta) = \chi_0(i).$$

If $i < |F|$, then $F_\alpha(i) = F_\beta(i)$ and, since β is in U_β and $F < \beta$,

$$c(F_\alpha(i), \beta) = c(F_\beta(i), \beta) = \chi_0(i) = 1.$$

If $i \geq |F|$, then $\chi_0(i) = \chi(i - |F|)$ if $i - |F| < l$ and $\chi_0(i) = \chi(i - |F| + 1)$ if $i - |F| \geq l$. If $i - |F| < l$, then

$$c(F_\alpha(i), \beta) = c(a(i - |F|), \beta) = \chi(i - |F|) = \chi_0(i)$$

and if $i - |F| \geq l$, then

$$c(F_\alpha(i), \beta) = c(a(i - |F| + 1), \beta) = \chi(i - |F| + 1) = \chi_0(i).$$

This finishes the proof. $\square$

Corollary 7.8. *For every X , $\tau[X]$ is hereditarily Lindelöf.*

Proof. If not, then $(X, \tau[X])$ would contain an uncountable discrete subspace. It is then possible to find disjoint $Y, Z \subseteq X$ such that $(Y, \tau[Y])$ and $(Z, \tau[Z])$ each contain uncountable discrete subspaces. But any function from a discrete space to another discrete space is continuous and we can then easily contradict Theorem 7.7. $\square$

Kunen has shown, however, that under $\text{MA}_{\aleph_1}$ every L space contains an uncountable discrete subspace in one of its finite powers [13]. As might be expected, this happens at the first possible instance in our example.

Theorem 7.9. *For every uncountable $X \subseteq \omega_1$, $\tau[X]^2$ contains an uncountable discrete subspace.*

Proof. This essentially follows from the following observation which is of independent interest.

Proposition 7.10. *The tree*

$$T(o) = \{o(\cdot, \beta) \upharpoonright \alpha : \alpha \leq \beta < \omega_1\}$$

is Aronszajn.

Proof. Theorem 5.3 implies that $T(o)$ has no uncountable branches. It suffices to prove that all levels of $T(o)$ are countable. For this, it is sufficient to show that $T(\varrho_1)$ and $T(\mu)$ have countable levels. In the case of $T(\varrho_1)$, this is a trivial consequence of Fact 5. In order to see that $T(\mu)$ has countable levels, let $\alpha < \omega_1$ be given. Let $\beta < \omega_1$ be greater than α . Using the compactness of $\alpha + 1$, select a finite set $F_\beta \subseteq \alpha + 1$ containing 0 and α so that if $\gamma_0 < \gamma$ are consecutive elements of F_β , then

$$L(\gamma, \beta) < L(\xi, \gamma)$$

whenever $\gamma_0 < \xi < \gamma$. It suffices to show that if

$$\mu(\cdot, \beta) \upharpoonright F_\beta = \mu(\cdot, \beta') \upharpoonright F_{\beta'},$$

then

$$\mu(\cdot, \beta) \upharpoonright \alpha = L(\cdot, \beta') \upharpoonright \alpha.$$

To see this, let $\xi < \alpha$ be arbitrary and pick $\gamma_0 < \gamma$ in F_β so that $\gamma_0 < \xi \leq \gamma$. If $\xi = \gamma$, then $\mu(\xi, \beta) = \mu(\xi, \beta')$ by assumption. Otherwise,

$$\mu(\xi, \beta) = \mu(\xi, \gamma) \cup \mu(\gamma, \beta) = \mu(\xi, \gamma) \cup \mu(\gamma, \beta') = \mu(\xi, \beta').$$

□

To finish the proof of Theorem 7.9, select a sequence $(\beta_\xi^0, \beta_\xi^1)$ indexes by an uncountable set $\Xi \subseteq \omega_1$ such that for each ξ in Ξ the following conditions are satisfied:

- (1) $\beta_\xi^0 < \beta_\xi^1$ are both elements of X .
- (2) Both β_ξ^0 and β_ξ^1 are greater than ξ .
- (3) If $\eta < \xi$ is in Ξ , then $\beta_\eta^1 < \xi$.
- (4) $o(\cdot, \beta_\xi^0) \upharpoonright \xi = o(\beta_\xi^1) \upharpoonright \xi$.

This is possible since $T(o)$ is Aronszajn. Now consider the clopen neighborhoods

$$U_\xi = (W_{\beta_\xi^1} \setminus W_{\beta_\xi^0}) \times W_{\beta_\xi^1}$$

of $(\beta_\xi^0, \beta_\xi^1)$ for ξ in Ξ . If $\eta < \xi$ are in Ξ , then $(\beta_\eta^0, \beta_\eta^1)$ is not in U_ξ . If $\xi < \eta$ are in Ξ , then

$$o(\beta_\xi^1, \beta_\eta^0) = o(\beta_\xi^1, \beta_\eta^1).$$

It follows that β_η^0 is in $W_{\beta_\xi^1}$ iff β_β^1 is. This means that $(\beta_\eta^0, \beta_\eta^1)$ is not in U_ξ . Hence $(\beta_\eta^0, \beta_\eta^1)$ is in U_ξ iff $\eta = \xi$ and therefore

$$\{(\beta_\xi^0, \beta_\xi^1) : \xi \in \Xi\}$$

is an uncountable discrete subspace of $\tau[X]^2$ as desired. □

The proof of Theorem 7.3 can be used to prove the following.

Theorem 7.11. *For every $X \subseteq \omega_1$, no uncountable subspace of $\tau[X]$ is first countable.*

Proof. The proof of Theorem 7.3 actually shows that if an L space is not first countable, then there is a c.c.c. forcing which destroys it. It is easily checked that the any uncountable subspace of $\tau[X]$ is a c.c.c. indestructible L space. □

We also have the following consequence of Proposition 7.10.

Theorem 7.12. *For every X , if f is a continuous function from $\tau[X]$ into $[0, 1]$, the range of f is countable.*

Proof. For each rational q in $[0, 1]$, let U_q be the f -preimage of $[0, q)$. Applying the hereditary Lindelöf property of $\tau[X]$, find countable $\mathcal{U}_q$ consisting of basic clopen sets such that $U_q = \cup \mathcal{U}_q$. Now let δ be an ordinal larger than any ordinal mentioned in the definition of an element of some $\mathcal{U}_q$. Applying Proposition 7.10, pick β_0 and β_1 in X such that $c(\cdot, \beta_0) \upharpoonright \delta = c(\cdot, \beta_1) \upharpoonright \delta$. It suffices to show that $f(\beta_0) = f(\beta_1)$. If not, pick a rational such that $[0, q)$ contains exactly one of

$f(\beta_0), f(\beta_1)$. But then there is an element of $\mathcal{U}_q$ which contains exactly one of β_0, β_1 . This means that for some $\alpha < \delta$,

$$c(\alpha, \beta_0) \neq c(\alpha, \beta_1),$$

contradicting the definition of δ and our choice of β_0, β_1 . $\square$

Theorem 7.13. *1.13 Any basis for the uncountable regular Hausdorff spaces must have cardinality at least $\aleph_2$.*

Proof. The first conclusion follows from Theorem 7.7 and the observation that there is an almost disjoint family of uncountable subsets of ω_1 of cardinality $\aleph_2$. $\square$

REFERENCES

- [1] Noga Alon and Joel H. Spencer. *The probabilistic method*. Wiley-Interscience Series in Discrete Mathematics and Optimization. Wiley-Interscience [John Wiley & Sons], New York, second edition, 2000.
- [2] James E. Baumgartner. All $\aleph_1$ -dense sets of reals can be isomorphic. *Fund. Math.*, 79(2):101–106, 1973.
- [3] M. M. Day. Oriented systems. *Duke Math J.*, 11:201–229, 1944.
- [4] Paul Erdős and Andras Hajnal. Embedding theorems for graph establishing negative partition relations. *Periodica Mathematica Hung.*, 9:205–230, 1978.
- [5] David H. Fremlin. Problems list.
- [6] David H. Fremlin. *Consequences Of Martin's Axiom*. Cambridge University Press, 1984.
- [7] Gary Gruenhage. Cosmicity of metrizable spaces. *Trans. Amer. Math. Soc.*, 313(1):301–315, 1989.
- [8] Gary Gruenhage. Perfectly normal compacta, cosmic spaces, and some partition problems. In *Open problems in topology*, pages 85–95. North-Holland, Amsterdam, 1990.
- [9] A. Hajnal and I. Juhász. On hereditarily α -Lindelöf and hereditarily α -separable spaces. *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.*, 11:115–124, 1968.
- [10] T. Jech. *Multiple forcing*, volume 88 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1986.
- [11] Thomas Jech. *Set theory*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, second edition, 1997.
- [12] I. Juhász. A survey of S - and L -spaces. In *Topology, Vol. II (Proc. Fourth Colloq., Budapest, 1978)*, volume 23 of *Colloq. Math. Soc. János Bolyai*, pages 675–688. North-Holland, Amsterdam, 1980.
- [13] Kenneth Kunen. Strong S and L spaces under MA . In *Set-theoretic topology (Papers, Inst. Medicine and Math., Ohio Univ., Athens, Ohio, 1975-1976)*, pages 265–268. Academic Press, New York, 1977.
- [14] Kenneth Kunen. *An introduction to independence proofs*, volume 102 of *Studies in Logic and the Foundations of Mathematics*. North-Holland, 1983.
- [15] C. Kuratowski. Sur le problème des courbes gauches en topologie. *Fund. Math.*, 15:271–283, 1930.

- [16] Dj. Kurepa. Ensembles ordonnés et ramifiés. *Publ. Math. Univ. Belgrade*, 4:1–138, 1935.
- [17] Paul Larson and Stevo Todorčević. Katětov’s problem. *Trans. Amer. Math. Soc.*, 354:1783–1792, 2002.
- [18] Justin Tatch Moore. A five element basis for the uncountable linear orders. Preprint, February 2004.
- [19] Judy Roitman. Adding a random or a cohen real. *Fund. Math.*, 103:47–60, 1979.
- [20] Judy Roitman. Basic S and L . In *Handbook of set-theoretic topology*, pages 295–326. North-Holland, Amsterdam, 1984.
- [21] Mary Ellen Rudin. S and L spaces. In *Surveys in general topology*, pages 431–444. Academic Press, New York, 1980.
- [22] Saharon Shelah. Decomposing uncountable squares to countably many chains. *J. Combinatorial Theory Ser. A*, 21(1):110–114, 1976.
- [23] J. R. Shoenfield. The problem of predicativity. In *Essays on the foundations of mathematics*, pages 132–139. Magnes Press, 1961.
- [24] W. Sierpiński. Sur un problème concernant les types de dimensions. *Fundamenta Mathematicae*, 19:65–71, 1932.
- [25] M. Suslin. Problème 3. *Fund. Math.*, 1:223, 1920.
- [26] Zoltán Szentmiklóssy. S spaces and L spaces under Martin’s Axiom. In *Topology*, volume 23 of *Coll. Math. Soc. Janos Bolyai*, pages 1139–1145. North-Holland, 1980. Fourth Colloq., Budapest 1978.
- [27] Stevo Todorčević. Forcing positive partition relations. *Trans. Amer. Math. Soc.*, 280(2):703–720, 1983.
- [28] Stevo Todorčević. Basis problems in combinatorial set theory. In *Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998)*, number Extra Vol. II, pages 43–52, 1998.
- [29] Stevo Todorčević. Compact subsets of the first Baire class. *J. Amer. Math. Soc.*, 12(4):1179–1212, 1999.
- [30] Stevo Todorčević. Partitioning pairs of countable ordinals. *Acta Math.*, 159(3–4):261–294, 1987.
- [31] Stevo Todorčević. *Partition Problems In Topology*. Amer. Math. Soc., 1989.
- [32] Stevo Todorčević. A classification of transitive relations on ω_1 . *Proc. London Math. Soc. (3)*, 73(3):501–533, 1996.
- [33] Stevo Todorčević. Coherent sequences. In *Handbook of Set Theory*. North-Holland, (in preparation).
- [34] J.W. Tukey. Convergence and uniformity in topology. *Princeton Univ. Press*, 1940.
- [35] Peter Vojtáš. Generalized Galois-Tukey-connections between explicit relations on classical objects of real analysis. In *Set theory of the reals (Ramat Gan, 1991)*, pages 619–643. Bar-Ilan Univ., Ramat Gan, 1993.
- [36] W. Hugh Woodin. *The Axiom of Determinacy, Forcing Axioms, and the Non-stationary Ideal*. Logic and its Applications. de Gruyter, 1999.
- [37] Phillip Zenor. Hereditary $\mathfrak{m}$ -separability and the hereditary $\mathfrak{m}$ -Lindelöf property in product spaces and function spaces. *Fund. Math.*, 106(3):175–180, 1980.

DEPARTMENT OF MATHEMATICS, BOISE STATE UNIVERSITY, BOISE, ID 83724
E-mail address: justin@math.boisestate.edu